

- 13 **Eves ND**, Petersen SR, Jones RL. The effects of helium-oxygen (HE-OX) and helium-hyperoxia (HE-HOX) during graded exercise with the self-contained breathing apparatus. *Can J Appl Physiol* 2001;**2**:477.
- 14 **Forster HV**, Erickson BK, Lowry TF, *et al.* Effect of helium-induced ventilatory unloading on breathing and diaphragm EMG in awake ponies. *J Appl Physiol* 1994;**77**:452–62.
- 15 **Christopherson SK**, Hlastala MP. Pulmonary gas exchange during altered density gas breathing. *J Appl Physiol* 1982;**52**:221–5.
- 16 **Woods SL**. Monitoring pulmonary artery pressures. *Am J Nurs* 1976;**76**:1765–71.
- 17 **Nemery B**, Nullens W, Veriter C, *et al.* Effects of gas density on pulmonary gas exchange of normal man at rest and during exercise. *Pflügers Arch* 1983;**397**:57–61.
- 18 **Krishnan BS**, Clemens RE, Zintel TA, *et al.* Ventilatory response to helium-oxygen breathing during exercise: effect of airway anesthesia. *J Appl Physiol* 1997;**83**:82–8.
- 19 **Dempsey JA**, Hanson PG, Henderson KS. Exercise-induced arterial hypoxaemia in healthy human subjects at sea level. *J Physiol* 1984;**355**:161–75.
- 20 **Young AJ**, Cymerman A, Pandolf KB. Differentiated ratings of perceived exertion are influenced by high altitude exposure. *Med Sci Sports Exerc* 1982;**14**:223–8.
- 21 **Dressendorfer RH**, Wade CE, Bernauer EM. Combined effects of breathing resistance and hyperoxia on aerobic work tolerance. *J Appl Physiol* 1977;**42**:444–8.
- 22 **Eves ND**, Petersen SR, Haykowsky MJ, *et al.* Helium-hyperoxia, exercise and respiratory mechanics in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2006.
- 23 **Weston SB**, Gabbett TJ. Reproducibility of ventilation of thresholds in trained cyclists during ramp cycle exercise. *J Sci Med Sport* 2001;**4**:357–66.
- 24 **Noble BJ**, Borg GA, Jacobs I, *et al.* A category-ratio perceived exertion scale: relationship to blood and muscle lactates and heart rate. *Med Sci Sports Exerc* 1983;**15**:523–8.
- 25 **Babb TG**. Breathing He-O<sub>2</sub> increases ventilation but does not decrease the work of breathing during exercise. *Am J Respir Crit Care Med* 2001;**163**:1128–34.
- 26 **Richardson RS**, Sheldon J, Poole DC, *et al.* Evidence of skeletal muscle metabolic reserve during whole body exercise in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1999;**159**:881–5.
- 27 **Puente-Maestu L**, Garcia de PJ, Martinez-Abad Y, *et al.* Dyspnea, ventilatory pattern, and changes in dynamic hyperinflation related to the intensity of constant work rate exercise in COPD. *Chest* 2005;**128**:651–6.
- 28 **Debinski W**, Klossowski M, Gembicka D. Effect of breathing of a helium-oxygen mixture on adaptation to effort in humans during high-altitude hypoxia. *Acta Physiol Pol* 1986;**37**:32–40.
- 29 **Aaron EA**, Seow KC, Johnson BD, *et al.* Oxygen cost of exercise hyperpnea: implications for performance. *J Appl Physiol* 1992;**72**:1818–25.
- 30 **Dempsey JA**, Wagner PD. Exercise-induced arterial hypoxemia. *J Appl Physiol* 1999;**87**:1997–2006.
- 31 **Ansley L**. PhD thesis 2003.
- 32 **Laplaid D**, Hug F, Grelot L. Reproducibility of eight lower limb muscles activity level in the course of an incremental pedaling exercise. *J Electromyogr Kinesiol* 2006;**16**:158–66.
- 33 **Kuipers H**, Verstappen FT, Keizer HA, *et al.* Variability of aerobic performance in the laboratory and its physiologic correlates. *Int J Sports Med* 1985;**6**:197–201.

## COMMENTARY

This paper provides relevant data on furthering our understanding of cardiovascular dynamics during exercise, which may well impact on sporting performance and have clinical significance.

**A M Hunter**

University of Stirling, Stirling, UK; a.m.hunter1@stir.ac.uk

## EDITORIAL BOARD MEMBER

### Mark Batt

**P**rofessor Mark Batt, BSc MB BChir MRCP DM FFSEM FACS, is a consultant in sport and exercise medicine at The Centre for Sports Medicine, University Hospitals NHS Trust, Nottingham. He has a busy NHS practice and is physician for The English Institute of Sport.

He graduated from Cambridge University Medical School in 1984 and trained in family medicine. He obtained a diploma in sports medicine from the University of London in 1991 and completed a fellowship in sports medicine at the University of California, Davis (UCD) in 1993. The next two years were spent as a faculty member in Family Medicine at UCD and as a team physician at the University of California, Berkeley.

Since 1995, he has been in Nottingham as a consultant/senior lecturer in sport and exercise medicine at the Queens Medical Centre: appointed special professor in 2004. He is currently clinical director for trauma and orthopaedics. He acts as clinical advisor for the Nottingham MSc/diploma courses in sports medicine. He serves or served as a consultant for The England and Wales Cricket Board, The Rugby Football League, British Gymnastics, The English Institute of Sport and The Wimbledon Tennis Championships. He is vice-chairman of the Faculty of Sport and Exercise Medicine and chaired the work-group that produced the case for sport and exercise medicine as a specialty of medicine. He is chairman of the newly created Specialist Advisory Committee in Sport and Exercise Medicine.

His research interests include: overuse injuries, particularly groin, low back, lower leg pain (shin splints and stress fractures); tendon disease; and exercise in the workplace.

He is married with two children. He enjoys a variety of sports, outdoor pursuits and gardening, none of which he does tremendously well!



**Figure 1** Mark Batt.

doi: 10.1136/bjism.2006.031880